

VACCINES

Hepatitis A

WHO

- Gay, bisexual, or other men who have sex with men.
- Trans women
- Anyone living with HIV, Hepatitis B or Hepatitis C

WHAT

- Two doses, at least 6 months apart
- Combined hepatitis A and B vaccine available but schedules will be different

Hepatitis B

WHO

- Gay, bisexual, or other men who have sex with men
- Trans women
- Anyone living with HIV, Hepatitis B or Hepatitis C
- People whose partner has Hepatitis B
- Sex workers and those who have sex with sex workers

WHAT

- Three doses with or without a booster
- Schedules vary but generally given within 3 months
- Can be given within 3 weeks if faster protection required
- Combined hepatitis A and B vaccine available but schedules may be different

HPV

WHO

- Gay, bisexual, or other men who have sex with men up to aged 45 and under
- Trans women aged 45 and under

WHAT

- One dose if under 25
- Two doses, at least 6 months apart, if 25-45

MPOX

VACCINE NUMBERS ARE LIMITED AND CAN ONLY BE GIVEN IF CERTAIN CRITERIA ARE MET

WHO

- Gay, bisexual, or other men who have sex with men who have multiple sexual partners, attend sex on premises venues, or engage in group sex

WHAT

- Two doses, at least 1 month apart

4CMenB (Gonorrhoea)

VACCINE NUMBERS ARE LIMITED - ONLY GIVEN IF CERTAIN CRITERIA ARE MET

WHO

- Gay, bisexual, or other men who have sex with men who have multiple sexual partners, attend sex on premises venues, or engage in group sex

WHAT

- Two doses, at least one month apart
- 30-40% effectiveness

HIV PREVENTION

Post Exposure Prophylaxis (PEPSE)

WHO

- People who are deemed to be HIGH risk of acquiring HIV through sexual contact
- This includes certain types of unprotected sex with people of unknown status or known HIV positive with a detectable viral load

WHAT

- Three tablets daily for 28 days
- Started within 72 hours after potential exposure

Pre-Exposure Prophylaxis (PrEP)

WHO

- Gay, bisexual, or other men who have sex with men who engage in unprotected sex
- Trans women
- Other people who feel they may be at increased risk of HIV

WHAT

- Generally taken as a once daily tablet
- May be taken "event based" depending on type of sex

STI PREVENTION

DoxyPEP

WHO

- Gay, bisexual, and other men who have sex with men
- Trans women

WHAT

- Single dose of doxycycline to be taken within 72 hours after unprotected sex
- Reduced risk of syphilis and chlamydia

Condoms

WHO

- Anyone

WHAT

- External and Internal condoms available
- Protection against STIs and pregnancy